

A Survey for Large Separation Lensed FIRST Quasars, II. Magnification Bias and Redshift Distribution

Eran O. Ofek^{★1,2}, Hans-Walter Rix², Dan Maoz¹, Francisco Prada³

¹ *School of Physics and Astronomy and Wise Observatory, Tel Aviv University, Tel Aviv 69978, Israel*

² *Max-Planck-Institut für Astronomie, Königstuhl 17, D-69117 Heidelberg, Germany*

³ *Isaac Newton Group, Apartado de Correos 368, Santa Cruz de La Palma, 38750 Tenerife, Canary Islands, Spain*

Accepted ? Received ? in original form ?

ABSTRACT

The statistics of large-separation gravitational lensing are a powerful tool to probe mass distributions on the scale of galaxy clusters. In this paper we refine the analysis of our survey for large-separation ($> 5''$) lensed FIRST quasars (Ofek et al. 2001) by estimating the magnification bias and the source redshift distribution. Finding no large separation lens among 8000 likely quasars in that sample, implies an upper bound on the lensed fraction of 3.7×10^{-4} at 95% CL. From a published deep 1.4 GHz radio survey of the Hubble Deep Field, and corresponding optical searches for faint quasars, we calculate a lower limit to the ‘double flux magnification bias’ affecting our radio-optically selected sample, of $B \geq 1.1$. From the four-colour information in the SDSS Early Data Release, we calculate the photometric redshift distribution of a sample of FIRST quasar candidates and compare it with the redshift distribution from the FIRST Bright Quasar Survey. We find that the median redshift of the quasars in our sample is about 1.4. With these new results, we find that for all plausible cosmologies, the absence of lensed quasars in our survey is consistent with a model based on an empirical, non-evolving, cluster mass function, where clusters are represented by singular isothermal spheres. On the other hand, comparison of our results to the lensing predictions of published N -body-ray-tracing simulations (Wambsganss et al. 1995, 1998) rejects the COBE normalised $\Omega_0 = 1$ CDM model at 99.9% confidence.

Key words: cosmology: gravitational lensing – galaxies: clusters: general – quasars: general

1 INTRODUCTION

The statistics of gravitational lensing can provide a powerful probe of the geometry and the mass content of the universe out to large redshifts (e.g. Refsdal 1964; Press & Gunn 1973; Turner, Ostriker, & Gott 1984). Lensed quasar statistics in the angular image separation ranges expected from galaxy-mass lenses have been probed by several surveys (e.g., Maoz et al. 1993). However, systematic searches for lensed quasars with multiple image separations of $5'' \lesssim \Delta\theta \lesssim 30''$ expected from groups or clusters of galaxies have been carried out only recently (e.g., Maoz et al. 1997; Marlow et al. 1998; Phillips et al. 2001a; Phillips, Browne, & Wilkinson 2001; Ofek et al. 2001; Keeton & Madau 2001; Phillips et al. 2001b).

The incidence of large-separation gravitational lensing depends on several factors, including the cosmology, the present-day cluster mass function and its redshift evolution,

and most importantly, the mass profile and the substructure within galaxy clusters (e.g., Bartelmann, Steinmetz, & Weiss 1995). There are several approaches for predicting the optical depth to large-separation lensing in a given model scenario:

(i) Ray tracing through N -body simulations (e.g., Cen et al. 1994; Wambsganss et al. 1995; Wambsganss, Cen, & Ostriker 1998). This accounts for the complicated large-scale, multi-plane lens structure, and for the substructure in clusters of galaxies. Wambsganss et al. (1995; 1998) predict an optical depth for lensing with $\Delta\theta > 5''$ of about 3×10^{-3} , for a $z \approx 3$ source in an $\Omega_0 = 1$ Cold Dark Matter (CDM) cosmology. The problem of substructure, in particular, was analysed by Bartelmann & Weiss (1994), and Bartelmann et al. (1995). They found that numerically modeled clusters with substructure are about two orders of magnitude more efficient in producing highly magnified images (e.g., long arcs) than spherically symmetric clusters.

(ii) Taking the observed cluster mass function and adopting a smooth and symmetric mass profile to calculate

[★] e-mail: eran@wise.tau.ac.il

the optical depth (e.g., Maoz et al. 1997; Ofek et al. 2001). Maoz et al. (1997) used the NFW profile (Navarro, Frenk, & White 1996, 1997) to show that the cluster lensing optical depth can be in the range $\tau(\Delta\theta > 5'') \approx 10^{-5}$ to 10^{-2} , where τ depends mostly on the adopted mass-concentration. In Ofek et al. 2001 (hereafter Paper I), we estimated the optical depth for singular isothermal sphere (SIS) lenses using the non-evolving Girardi et al. (1998) cluster mass function and found $\tau(\Delta\theta > 5'') \approx 3 \times 10^{-4}$, for an $\Omega_0 = 0.3$, $\Omega_\Lambda = 0.7$ cosmology and a source redshift of $z_{\text{source}} = 1$.

(iii) Modifying approach (ii) by assuming the cluster mass function is given by the Press-Schechter formalism (Press & Schechter 1974), that by itself depends on cosmology. For example, Li & Ostriker (2002) have followed this approach, and assumed a generalized NFW density profile and $z_{\text{source}} = 1$. Their predictions for the optical depth span values in an even larger range, $7 \times 10^{-7} \lesssim \tau(\Delta\theta > 5'') \lesssim 10^{-3}$, depending in part on the details of the adopted mass profile. The impact of the cosmological model on the cross sections may seem surprisingly large, about one or two orders of magnitudes. This dependence enters through the differing redshift evolutions of the cluster mass function. Other examples of the Press-Schechter approach to cluster lensing statistics are Narayan & White (1988), Flores & Primack (1996), Wyithe, Turner, & Spergel (2001), and Sarbu, Rusin, & Ma (2001).

In light of these wide-ranging predictions, empirical constraints seem in order. Without any confirmed large-separation quasar lenses, we can only place an upper limit on the cross section which, in turn, can reject part of the mass-profile and cosmology parameter space.

In Paper I we described a survey for large separation gravitational lensing among quasars drawn from the FIRST radio catalog. In that survey, we selected radio sources from the FIRST catalogue (1999, July 21 version; Becker, White, & Helfand 1994; White et al. 1997) with optical counterparts in the APM catalog (McMahon, & Irwin 1992), with positional coincidence better than $2.''5$ in both the O and E bands, with colour index $O - E < 2$ mag, and that are point-like on at least one of the O or E POSS-I/UKST plates, as determined by the APM classification algorithm. There are 12,576 quasar candidates passing the above criteria. Based on the initial results from the FIRST Bright Quasar Survey (FBQS; Gregg et al. 1996; White et al. 2000), we estimated in Paper I that there are about 9100 quasars in our sample out of the 12,576 candidates.

However, with the more recently published results of the FBQS (Becker et al. 2001), which extends the spectroscopic follow-up from 17.8 to 19 mag, we can revise the fraction of quasars among the sources fainter than $E = 17.8$ mag in our sample from 90% to 80%, or about 8000 quasars instead of 9100 quasars. Specifically, Equation 1 in Paper I can be revised to:

$$F_{qso} = \begin{cases} -0.9389 + 0.046E + 0.0027E^2, & E < 18 \\ 0.8, & E \geq 18 \end{cases} \quad (1)$$

Follow-up observations of all pairs of radio-optical quasar candidates with $5'' < \Delta\theta < 30''$ showed that none of the 15 candidate pairs are lensed quasars (for details see Paper I). For 8000 quasars in the sample, this implies an upper limit of 3.7×10^{-4} (95% Poisson statistics confidence level; CL) on the lensing fraction in this survey.

However, in order to constrain the mass function and mass profile of clusters from these results, some further properties of our survey need to be known: (i) the double flux magnification bias of our survey, resulting from the radio and optical flux limits involved in the sample selection. This magnification bias reflects the over- (or under) representation of lensed objects in our survey, compared to purely geometrical cross-section calculations; and (ii) the redshift distribution of quasars (i.e., the potential sources) in our sample, which determines the effective pathlength of our lens search.

The purpose of the present paper is to improve those two important constraints and provide a more reliable upper limit on the observed optical depth to large separation lensing.

2 THE SURVEY DOUBLE FLUX MAGNIFICATION BIAS

As described in Paper I, the magnification bias depends strongly on the faint-end number counts (in the radio and optical) of quasars with similar selection criteria to those used in our survey. However, there are presently no large-area very deep radio-optical surveys. The best available data set is the combination of radio and optical surveys in the Hubble Deep Field North (HDF-N) region: a deep 1.4 GHz radio survey conducted with the VLA by Richards (2000). This survey has a flux limit (5σ) of 40 μJy in the central $8'$, degrading to about 300 μJy at $18'$ from the field centre. In the optical, a 1 deg^2 region around the HDF-N was searched for quasars down to $B = 22$ mag using multi-colour selection (Liu et al. 1999; Vanden Berk et al. 2000). Liu et al. required somewhat more restrictive color criteria than ours: $B - R < 0.8$, or $U - B \leq -0.4$ and $B - R \leq 1.1$. Impey & Petry (2001) matched the Richards (2000) radio sources with the optically detected quasars in this field, and identified three radio-optical quasars within the colour range, $0.46 < B - R < 0.75$, that match our survey criteria. With sufficient lensing magnification these quasars would have been detected by our survey. These data on the HDF-N constitute faint-end counts that can be combined with our FIRST-APM candidates, to find the double flux magnification bias.

The HDF implies radio counts of 190^{+180}_{-100} radio-optical sources per deg^2 per mJy, at 0.12 mJy in the 1.4 GHz-band, and optical counts of $3.5^{+8.1}_{-2.9}$ radio-optical sources per deg^2 per mag at $O = 20.3$ magnitude. The new radio number-count point includes three radio-optical quasars with optical counterpart brighter than $B = 22$, and the optical point is based on one faint (fainter than the APM 100% completeness limit of 19.75; Caretta et al. 2000) radio-optical quasar with radio counterpart brighter than the Richards (2000) survey limit, as given above. Figure 1 shows the differential number counts in the 1.4 GHz-band for the Paper I sample quasars (circles), and for the deep radio-optical search for quasars described above (single square). The dashed line is the interpolated number-counts in the 0.1-1 mJy range. Figure 2 is the same as Figure 1, but for the optical O -band.

Although it is reasonable to assume that there is a large number of radio-faint (fainter than 0.1 mJy) quasars consti-




Figure 1. Differential number counts in the 1.4 GHz-band for our sample quasars (circles), and one square representing the deep radio-optical search for quasars described above. The dashed line is the interpolated number-counts in the 0.1-1 mJy range.




Figure 2. Same as Figure 1, but for the O-band

tuting the radio-quiet quasar population, we only have direct observational constraints on the number counts down to 0.1 mJy (Figures 1-2). Therefore, our estimate for the double-flux magnification bias will only be a lower limit. Furthermore, due to the small area of the HDF-N deep radio observations, the number counts at the intermediate flux levels (between FIRST and HDF-N) are not measured directly, and we simply interpolate the number count distribution over these gaps.

Drawing on this new information we have calculated the double flux magnification bias in the FIRST-APM sample through a Monte-Carlo simulation, as follows.

We constructed a list of radio-optical quasars that mimics the observed number count distribution of the O-magnitude and of the 1.4 GHz flux shown in Figures 1-2. For quasars above the FIRST-APM flux limits we drew at random objects from our list of quasar candidates, corrected for the fraction F_{qso} of these objects that are quasars, and for the

incompleteness of the APM survey (Caretta et al. 2000), by replicating objects in the magnitude-range 19.5-21.5, until we had a list of about 8000 radio-optical sources. At the faint end we added objects with radio flux and optical magnitude drawing from a flux probability distribution obtained by interpolating the number counts in Figure 1. For each of these objects we then drew an O-magnitude from the distribution given in Figure 2. Note that the last step assumes that there is no correlation between the optical and radio properties of quasars, which is true at least for objects brighter than 1 mJy and $O \sim 21$ (Paper I). The number of randomly selected objects in this step was chosen by scaling the number of HDF-N faint sources by the ratio of the total area of the FIRST-APM survey to the area of the HDF-N faint-end number-count survey described above.

With the list described above, we are now in a position to estimate the double-flux magnification bias, assuming a SIS lens mass profile. Our FIRST-APM survey will detect a lensed quasar as such only if *both* lensed images (bright and faint) are brighter than the radio 1 mJy flux limit, *and* are brighter than the O-magnitude limit of 21.2, which is the weighted average magnitude limit of the APM (Caretta et al. 2000). The magnification/de-magnification of the faint image, A_- , in a SIS lens is given by: $\beta = 1/(1+A_-)$, where β is the impact parameter in units of the Einstein ring radius. By comparing the magnitude and flux of each object in the Monte-Carlo list with the survey's flux limits, we calculated for each object, i , the minimum magnification needed (or maximum de-magnification allowed), $A_{i,min}$, and the corresponding $\beta_{i,max}$, needed for the detection of an image in both the radio and optical bands in our survey (i.e., the smaller among the optical and radio values of $\beta_{i,max}$ was chosen).

The cross-section for strong lensing is defined as the area with $\beta \leq 1$, and the probability for a given impact parameter β scales like the area, β^2 . To estimate the double-flux magnification bias, B , we summed $\beta_{i,max}^2$ for all the objects in our Monte-Carlo list and divided the results by the number of quasars in our sample, i.e., 8000. The resulting ratio reflects the over- (or under) representation of lensed objects in our survey, compared to a purely geometrical cross-section calculation.

Using the above method, we obtain a lower limit of $B \gtrsim 1.1$. This limit changes to 0.9 if we take into account the 1σ Poisson statistics (lower confidence interval of the number counts), and to 1.3 if we assume that the optical searches for quasars described above are 50% incomplete (see Impey & Petry 2001) due to the different colour selection process used by Liu et al. (1999) and Vanden Berk et al. (2000). The $B > 0.9$ limit we give above assumes that the number-counts are almost flat from $O \sim 19$ to $O \sim 20$, and then go to zero. The actual value of B could be larger due to the probable existence of a yet-fainter radio-optical quasar population. Hence, we consider $B \geq 1.1$ to be a conservative limit. For comparison, in Paper I we roughly estimated B by assuming that the product of the optical and radio number-flux relations is a power-law with a cutoff at a flux that is 0.5 – 4 orders of magnitude lower than our survey's flux/magnitude limits. Based on this, we estimated B to be in the range of 1 – 40. Since the HDF-N data extend the depth of our optical source counts by about 1 mag, the lower limit on B found in this paper is near the low-end estimate

of Paper I, but is now on a firmer observational basis. The present method we have used to calculate the bias is superior to the Borgeest, Linde, & Refsdal (1991) approach we used in Paper I, since it takes into account the actual shapes of the separate optical and radio number count distributions instead of replacing them by single power-law distributions.

3 REDSHIFT DISTRIBUTION OF THE FIRST/APM SOURCES

From the first major public release of the Sloan Digital Sky Survey - Early Data Release (SDSS-EDR; York et al. 2000; Stoughton et al. 2002), we constructed a sample of 443 point-like SDSS objects that: were selected as quasar-candidates in the SDSS-EDR based on their colors; are found within $1.''2$ from a FIRST radio source; and are brighter than $g' = 21$ mag. Their magnitudes were corrected for Galactic extinction (Schlegel, Finkbeiner, & Davis 1998). All but two sources in this list have $g' - r' < 2$ mag, so they are good representatives of our survey. Based on the SDSS quasar selection criteria (e.g., Newberg & Yanny 1997; York et al. 2000; Schneider et al. 2002), and on the FBQS (Gregg et al. 1996; Helfand et al. 1998; Becker et al. 1998; White et al. 2000; Becker et al. 2001), we know that the fraction of quasars in this sample is $\gtrsim 75\%$.

From the quasar redshift-colour relation, given by Richards et al. (2001a, Table 3), we estimated the photometric redshifts for our 443-object sample, by a simple χ^2 minimization (Richards et al. 2001b) between the predicted and the observed SDSS colours. The photometric-redshift distribution of these objects is shown as bars in Figure 3. The dashed line shows the redshift distribution of quasars in the FBQS faint extension ($E < 19$ mag; Becker et al. 2001). The photometric-redshift distribution is not changed considerably if only FIRST sources with SDSS counterparts brighter than $g' = 19$ mag are taken. However, our photometric redshift estimate of quasars has a considerable scatter: 30% of the redshifts have errors, $\Delta z > 0.25$. To estimate the effect of this scatter on the photometric redshift distribution, we have used the 898 quasars with known redshifts and SDSS colours given by Richards et al. (2001a) to calculate the distribution of differences between our redshift prediction and the observed redshifts in each 0.25 redshift bin. We convolved this scatter distribution as a function of redshift with the FBQS faint-extension redshift distribution ($E < 19$ mag; Becker et al. 2001) and the result is shown as the dotted line in Figure 3. Ideally, we would expect that the photometric redshift distribution (bars) be similar to the redshift distribution of FBQS quasars convolved with the photometric-redshift scatter (dotted line). However, our photometric-redshift distribution has an excess of quasars at $z \sim 3$, relative to the FBQS faint-extension redshift distribution. The median redshift for the sample of FBQS faint-extension quasars is $\bar{z} = 1.2$, and the median redshift we get from the photometric redshift down to limiting magnitude $g' = 21$ is $\bar{z} = 1.4$. There are two possibilities to explain the discrepancy: (i) only a fraction ($\gtrsim 75\%$) of the sources in our sample are real quasars. Therefore, the non-quasar objects could contaminate the redshift distribution in some systematic way; (ii) the photometric-redshift procedure of Richards et al. (2001b) was developed based mostly on radio-quiet




Figure 3. The redshift distribution of FIRST quasar candidates down to $g' = 21$ mag (bars). The redshift distribution was calculated using a simple colour-redshift relation. The superposed vertical-lines on the bars, are the Poisson error-bars due to the finite sample size. The dashed line is the redshift distribution of FBQS faint-extension ($E < 19$ mag) quasars. The dotted line is the same distribution, but convolved with the photometric redshift scatter, as a function of redshift.

quasars, but our survey contains mainly radio-loud quasars. It is known that there are some optical spectral differences between radio-loud and radio-quiet quasars (e.g., Brotherton et al. 2001). Such differences could bias the Richards et al. (2001b) photometric-redshift procedure. Therefore we adopt the FBQS faint-extension redshift distribution for our calculations. This is a conservative choice, as the lower median redshift implies a smaller path length for lensing.

4 DISCUSSION

In the following sections we use these two new pieces of information, $B \geq 1.1$ and the redshift distributions, to estimate the optical-depth to lensing in a simple SIS model, and to compare our results with $\Omega_0 = 1$ CDM N -body ray-tracing simulations.

4.1 A SIS model

To get some simple estimates of the lensing probability in our survey, we calculated the cross section as a function of redshift, assuming SIS lenses and the Girardi et al. (1998) mass function of groups and clusters with no evolution. We found in Paper I that the effective dimensionless density of lenses (as defined by Turner et al. 1984) is, $F = 16\pi^3 n_0 \left(\frac{c}{H_0}\right)^3 \left(\frac{\sigma_{\parallel}}{c}\right)^4 = 0.0156^{+0.012}_{-0.0067}$, where c is the speed of light, H_0 the Hubble constant, $\sigma_{\parallel}$ the line-of-sight velocity dispersion, and n_0 is the comoving number-density of lenses. The errors in F were obtained from Monte-Carlo simulations using the Girardi et al. (1998) mass function parameters and their errors (for details see Paper I). In Paper I we assumed a single representative redshift of $z = 1$ for all the quasars. With our new results presented above, we recalculate the total optical-depth for lensing, by integrating

the optical-depth over redshift, with the weighting function given by the FBQS faint-extension redshift probability density distribution (dashed-line in Figure 3). The optical depth τ_z , to a source at redshift z , is given by:

$$\tau_z = \int_0^z F(1+z_l)^3 \left(\frac{D_l D_{ls}}{D_s} \right)^2 \frac{1}{R_0} \frac{cdt}{dz_l} dz_l, \quad (2)$$

where D_l , D_{ls} , D_s are the angular-diameter distances for the observer-lens, lens-source and observer-source, respectively, z_l is the lens redshift, and $R_0 \equiv c/H_0$. The quantity cdt/dz_l is calculated in the Friedmann-Lemaître-Robertson-Walker geometry (e.g., Fukugita et al. 1992). The total weighted optical depth is given by:

$$\tau = \int_0^\infty P(z) \tau_z dz \quad (3)$$

where $P(z)$ is the normalized weighting function given by the FBQS quasar redshift distribution. We find

$$\tau = \begin{cases} 4.6^{+3.5}_{-2.0} \times 10^{-4}, & \Omega_0 = 0.3; \Omega_\Lambda = 0.7 \\ 2.4^{+1.8}_{-1.0} \times 10^{-4}, & \Omega_0 = 0.3; \Omega_\Lambda = 0.0 \\ 1.7^{+1.3}_{-0.7} \times 10^{-4}, & \Omega_0 = 1.0; \Omega_\Lambda = 0.0, \end{cases} \quad (4)$$

where the errors are the 1σ errors due to the uncertainty in F . This is ~ 1.7 times greater than estimated in Paper I, where $z = 1$ was assumed for all sources. Using the photometric-redshift distribution from Figure 3 (bars) instead of the FBQS faint-extension redshift distribution, the predicted τ would further increase by a factor of about 1.6. The difference is due to the excess of quasars with $z \sim 3$ in our photometric-redshift distribution, combined with the fact that the optical-depth increases with redshift. We conservatively adopt the optical-depth in Equation 4, as a lower limit on τ . These optical-depths need to be multiplied by the completeness factor that reflects the range of separations probed. As shown in Paper I, this completeness factor is about 0.66 for a SIS model. Finally, the lower limit on the magnification ($B \geq 1.1$) determines the expected probability for lensed quasars in our sample: $P_{lens} \geq 4.6^{+3.5}_{-2.0} \times 10^{-4} \times 0.66 \times 1.1 = 3.3^{+2.5}_{-1.5} \times 10^{-4}$ (for $\Omega_0 = 0.3$, $\Omega_\Lambda = 0.7$).

The observational upper bound on the lensed fraction of 3.7×10^{-4} (95% CL) from our survey (Paper I, and small revisions in this paper), is therefore consistent with this simple model (P_{lens}). Since the ($\Omega_0 = 0.3, \Omega_\Lambda = 0.0$) and ($\Omega_0 = 1.0, \Omega_\Lambda = 0.0$) models predict a lower optical depths, they too are consistent with the data. Within this simple model, for an $\Omega_0 = 0.3$, $\Omega_\Lambda = 0.7$ cosmology and a non-evolving Girardi et al. (1998) mass function, we can in turn set an upper limit to $F < 0.018$ with a 95% CL. Models incorporating evolution of the mass function will reduce the optical-depth, therefore increasing the upper limit on F .

4.2 Comparison with ray-tracing simulations

Wambsganss et al. (1995, 1998) performed ray tracing for $0 < z < 3$ through a three-dimensional mass distribution obtained from an $\Omega_0 = 1$ COBE normalized CDM N -body simulation. They found that the optical depth for strong lensing (i.e., for forming multiple images) with image separations $> 5''$ of sources at $z \sim 3$ is 3×10^{-3} . Our observations can test this model. Figure 2 in Wambsganss et al. (1995) describes the splitting probability for image pairs with separa-

tions greater than $5''$ and magnitude differences smaller than 1.5, as a function of the source redshift. To obtain the expected optical-depth for our survey, we integrate their splitting probability with a weight function given by the FBQS faint-extension source redshift distribution presented in Figure 3. We obtain $\tau = 1.14 \times 10^{-3}$ (or $\tau = 1.5 \times 10^{-3}$ using the photometric-redshift distribution). This optical depth needs to be multiplied by the completeness factor for the range of separations probed and the magnification bias of our survey. Wambsganss et al. (1995) also show (their Figure 3) the strong-lensing probability as a function of the image separation for various source redshifts. From this figure we estimate the incompleteness due to the separation range probed in our survey, $P(\Delta > 30'')/P(\Delta > 5'') \sim 0.25$ (This separation incompleteness is actually slightly redshift dependent, and changes from $\sim 30\%$ at $z_s = 1$ to $\sim 20\%$ at $z_s = 3$). Taking together τ , the separation incompleteness and the magnification bias we have found here, implies a probability for lensing in our survey $P_{CDM} \sim 1.14 \times 10^{-3} \times 0.75 \times 1.1 = 9.4 \times 10^{-4}$ based on the Wambsganss et al. simulations.

We note that the magnification bias we found in this paper is based on a SIS flux ratio distribution - an assumption that is not necessarily compatible with the Wambsganss et al. (1995, 1998) simulations. However, Figure 5 in Wambsganss et al. (1998) shows that the magnification probability found in their simulations is not very different from the $P(A) \sim A^{-3}$ expected from the SIS model we assumed in our magnification-bias calculation. Based on the calculation described above, 7.5 lenses are expected in our survey for an $\Omega_0 = 1$ CDM model. We can therefore reject the Wambsganss et al. (1995, 1998) model at 99.9% CL. We note that, since Figure 2 in Wambsganss et al. (1995) gives the splitting angle probability for image flux ratios smaller than 1.5 mag, our conclusion is conservative. Note, that the simulations of Wambsganss et al. use the COBE normalization, $\sigma_8 = 1.05$, which predicts an excess of high-mass clusters at the present epoch, and slower evolution of the mass function. In this sense, our results do not reject the $\Omega_0 = 1$ model *per se*, but the combination of the cosmology and the high normalization.

The results presented here constitute one of the tightest and well-defined limits on large separation lensing to date. Clearly, we have now reached a level where the non-detections are becoming interesting cosmological constraints. It is probable that actual cases of large-separation gravitationally lensed quasars will be found soon using one of the large area surveys (e.g., SDSS, 2DF; see Croom et al. 1998). However, in order to use such large separation lenses to constrain the cluster mass function and mass profile parameter space, we need to: (i) understand the selection effects and effectiveness of these surveys for large separation lensing; (ii) have a realistic estimate of the cross section for large separation lensing, taking into account the substructure of clusters. These points will be addressed in future papers.

To summarize our main results, we have derived a lower limit to the magnification bias in our survey, $B \geq 1.1$, and we have also found that the median photometric-redshift of quasars in our sample is $\bar{z}_{source} \sim 1.4$. Using the redshift distribution from the FBQS faint extension, we deter-

mine that $F < 0.018$ with 95% CL (assuming a SIS model and an $\Omega_0 = 0.3$, $\Omega_\Lambda = 0.7$ cosmology). Our non-detection of lensed FIRST quasars is consistent with expectations, if clusters can be represented by a non-evolving population of SIS masses with the mass function of the observed cluster population. If so, moderately larger surveys will discover the first examples of large separation quasar lensing. Our survey already has the ability to reject some models with concrete predictions, namely the $\Omega_0 = 1$ COBE-normalized CDM model, whose excess of power on large scales is well known.

ACKNOWLEDGMENTS

We thank Eric Richards for sending us a digital version of his deep radio catalogue of the HDF region, and an anonymous referee for useful comments. EEO wishes to thank the Max-Planck-Institut für Astronomie for its hospitality and financial support; the Deutscher Akademischer Austauschdienst for financial support; and Orly Gnat and Avishay Gal-Yam for fruitful discussions. This research has made use of the Sloan Digital Sky Survey. Funding for the creation and distribution of the SDSS Archive has been provided by the Alfred P. Sloan Foundation, the Participating Institutions, the National Aeronautics and Space Administration, the National Science Foundation, the U.S. Department of Energy, the Japanese Monbukagakusho, and the Max Planck Society. The SDSS Web site is <http://www.sdss.org/>.

REFERENCES

- Bartelmann, M., Weiss, A., 1994, *A&A*, 287, 1
 Bartelmann, M., Steinmetz, M., Weiss, A., 1995, *A&A*, 297, 1
 Becker, R.H., White, R.L., Helfand, D.J., 1994, in *Astronomical Data Analysis Software and Systems III*, ASP Conference Series, v. 61, eds. D. R. Crabtree, R. J. Hanisch, & J. Barnes, p. 165
 Becker, R.H. et al., 1998, *Am. Astron. Soc. Meeting*, 192, 11.01
 Becker, R.H. et al., 2001, *ApJS*, 135, 227
 Borgeest, U., von Linde, J., Refsdal, S., 1991, *A&A*, 251, L35
 Brotherton, M.S., Tran, H.D., Becker, R.H., Gregg, M.D., Laurent-Muehleisen, S.A., White, R.L., 2001, *ApJ*, 546, 775
 Caretta, C.A., Maia, M.A.G. & Willmer, C.N.A., 2000, *AJ*, 119, 524
 Cen, R., Gott, J.R., Ostriker, J.P., Turner, E.L., 1994, *ApJ*, 423, 1
 Flores, R.A., Primack, J.R., 1996, *ApJ*, 457, L5
 Fukugita, M., Futamase, T., Kasai, M., Turner, E.L., 1992, *ApJ*, 393, 3
 Croom, S.M., Shanks, T., Boyle, B.J., Smith, R.J., Miller, L., Loaring, N.S., 1998, *Evolution of Large-Scale Structure: From Recombination to Garching*, astro-ph/9810127
 Girardi, M., Borgani, S., Giuricin, G., Mardirossian, F., Mezzetti, M., 1998, *ApJ*, 506, 45
 Gregg, M.D., Becker, R.H., White, R.L., Helfand, D.J., McMahon, R.G., Hook, I.M., 1996, *AJ*, 112, 407
 Helfand, D.J., Yadigaroglu, I.A., Berger, R., Postman, M., White, R.L., Lauer, T.R., Oegerle, W.R., Becker, R.H., 1998, *Am. Astron. Soc. Meeting*, 193, 4002
 Impey, C., Petry, C., 2001, *ApJ*, 547, 117
 Keeton, C. R., Madau P., 2001, *ApJ*, 549, 25
 Li, L.-X., Ostriker, J.P., 2002, *ApJ*, 566, 652
 Liu, C.T., Petry, C.E., Impey, C.D., Foltz, C.B., 1999, *AJ*, 118, 1912
 Maoz, D., Bahcall, J.N., Schneider, D.P., Doxsey, R., Bahcall, N.A., Lahav, O. Yanny, B., 1993, *ApJ*, 402, 69
 Maoz, D., Rix, H.-W., Gal-Yam, A., Gould, A., 1997, *ApJ*, 486, 75
 Marlow D. R., McGuinness A. D., Browne I. W. A., Wilkinson P. N., Helbig P. J., 1998, *Large Scale Structure: Tracks and Traces*, eds. V. Mueller, S. Gottloeber, J.P. Muecket, J. Wambsganss, World Scientific, Singapore
 McMahon R.G., Irwin M.J., 1992, *Digitised Optical Sky Surveys*, Editors, H.T. MacGillivray, E.B. Thomson., Kluwer Acad. Press, Dordrecht
 Narayan, R., White, S.D.M., 1988, *MNRAS*, 231, 97
 Navarro, J., Frenk, C.S., White, S.D.M., 1996, *ApJ*, 462, 563
 Navarro, J., Frenk, C.S., White, S.D.M., 1997, *ApJ*, 490, 493
 Newberg, H.J., Yanny, B., 1997, *ApJS*, 113, 89
 Ofek, E.O., Maoz, D., Prada, F., Kolatt, T., Rix, H.-W., 2001, *MNRAS*, 324, 463
 Phillips, P.M., Browne, I.W.A., Wilkinson, P.N., Jackson, N.J., 2000, *New Cosmological Data and the Values of the Fundamental Parameters*, proceedings of IAU Symposium 201. Eds., A. Lasenby, Wilkinson, A., astro-ph/0011032
 Phillips, P.M., Browne, W.A., Wilkinson, P.N., 2001a, *MNRAS*, 321, 187
 Phillips, P.M., Browne, I.W.A., Jackson, N.J., Wilkinson, P.N., Mao, S., Rusin, D., Marlow, D.R., Snellen, I., Neeser, M., 2001b, *MNRAS*, 328, 1001
 Press, W.H., Gunn, J.E., 1973, *ApJ*, 185, 397
 Press, W.H., Schechter, P., 1974, *ApJ*, 187, 425
 Refsdal, S., 1964, *MNRAS*, 128, 307
 Richards, E.A., 2000, *ApJ*, 533, 611
 Richards, G.T. et al., 2001a, *AJ*, 121, 2308
 Richards, G.T. et al., 2001b, *AJ*, 122, 1151
 Sarbu, N., Rusin, D., Ma, C.-P., 2001, *ApJ*, 561, L147
 Schlegel, D.J., Finkbeiner, D.P., Davis, M., 1998, *ApJ*, 500, 525
 Schneider, D.P. et al., 2002, *AJ*, 123, 567
 Stoughton, C. et al., 2002, *AJ*, 123, 485
 Turner, E.L., Ostriker, J.P., Gott III, R., 1984, *ApJ*, 284, 1
 Vanden Berk, D.E., Stoughton, C., Crofts, A.P.S., Tytler, D., Kirkman, D., 2000, *AJ*, 119, 2571
 York, D.G. et al., 2000, *AJ*, 120, 1579
 Wambsganss, J., Cen, R., Ostriker, J.P., Turner, E.L., 1995, *Science*, 268, 274
 Wambsganss, J., Cen, R., Ostriker, J.P., 1998, *ApJ*, 494, 29
 White, R.L., Becker, R.H., Helfand, D.J., Gregg, M.D., 1997, *ApJ*, 475, 479
 White, R. L. et al., 2000, *ApJS*, 126, 133
 Wyithe, J.S.B., Turner, E.L., Spergel, D.N., 2001, *ApJ*, 555, 504